

Stable multiquark states with heavy quarks in a diquark model

Su Houn Lee^{1a} and Shigehiro Yasui^{2b}

¹ Institute of Physics and Applied Physics, Yonsei University, Seoul 120-749, Korea

² Institute of Particle and Nuclear Studies, High Energy Accelerator Research Organization (KEK), 1-1, Oho, Ibaraki 305-0801, Japan

Received: date / Revised version: date

Abstract. Based on the color-spin interaction in diquarks, we argue why some multiquark configurations could be stable against strong decay when heavy quarks are included. After showing the stability of previously discussed states we identify new possible stable states. These are the $T_{cb}^0(ud\bar{c}\bar{b})$ tetraquark, the $\Theta_{bs}(udus\bar{b})$ pentaquark and the $H_c(udusuc)$ dibaryon, and so forth.

PACS. 25.75.Nq Quark deconfinement, quark-gluon plasma production, and phase transitions – 12.39.Jh Nonrelativistic quark model – 14.80.-j Other particles (including hypothetical) – 14.20.Pt Dibaryons

1 Introduction

Multiquark configuration for the scalar nonet was first suggested by Jaffe in the framework of the MIT bag model [1]. Recently, the discussion on possible multiquark mesons has gained more ground as the recently discovered Z(4430) [2], Z(4051) and Z(4248) [3] seem to be composed of $c\bar{c}ud$ type of four quarks, as they decay into $\psi'\pi$ and $\chi_{c1}\pi$ respectively. The discussion on possibly multiquark configurations that are stable against strong decay is an old one starting from the H dibaryon ($udsuds$) [4], multiply charmed H dibaryon ($udsc\bar{c}\bar{c}$) or ($uudscc$) [5], charmed pentaquarks $\Theta_{cs}(udus\bar{c})$ [6, 7, 8, 9, 10, 11, 12] to tetraquarks $T_{cc}(ud\bar{c}\bar{c})$ [13, 14, 15]. While there are different models predicting the stability of these states, in the constituent quark model, the stability of possible stable multiquark configurations originate from the strong scalar diquark attraction [1, 16]. The strong attraction in the color antitriplet channel is an important ingredient in hadron spectroscopy and is also responsible for color superconductivity at high density [17]. In that sense, finding a stable multiquark configuration is also an important step toward understanding QCD at high density [18].

In this work, we introduce a simple model that can be used to analyze the stability of multiquark configuration against strong decay and show why some of the previously discussed states could indeed be stable. Then we discuss several new stable multiquark states, namely tetraquark, pentaquark and dibaryon containing charm and bottom quarks. The paper is organized as follows. In section II, we introduce our simple constituent quark model. Based on this model, we discuss possible stable multiquark states

and their measurable decay modes in section III. The summary is given in section IV.

2 A schematic model for hadron mass differences

It is a well known fact that the color-spin interaction [19] plays an important role in hadron mass differences. To illustrate the mechanism, we introduce the following simplified form for the color-spin interaction :

$$C_H \sum_{i>j} \mathbf{s}_i \cdot \mathbf{s}_j \frac{1}{m_i m_j}. \quad (1)$$

Here m_i and $\mathbf{s}_i$ are the mass and spin of the constituent quark i . The overall strength of the color-spin interaction is given as [20]

$$C_H = v_0 \boldsymbol{\lambda}_i \cdot \boldsymbol{\lambda}_j \langle \delta(r_{ij}) \rangle, \quad (2)$$

where v_0 is related to the coupling, the second part is the color factor and the third part is the strength of the wave function at zero separation. The color factor $\boldsymbol{\lambda}_i \cdot \boldsymbol{\lambda}_j$ would be $-8/3$ for diquarks in the color antitriplet channel and $-16/3$ for quark and anti-quark pair in the color singlet channel. It is a well known fact that the wave function effect can almost be treated as a constant [18, 22]. To illustrate this point, we assume the following constituent quark masses: $m_{u,d} = 300$ MeV, $m_s = 500$ MeV, $m_c = 1500$ MeV, and $m_b = 4700$ MeV.

Table 1 shows the mass differences between baryons that are sensitive to the color-spin interaction only. By fitting C_B to $M_\Delta - M_N$, we obtain $C_B/m_u^2 = 193$ MeV and find that the mass differences $M_\Sigma - M_\Lambda$, $M_{\Sigma_c} - M_{\Lambda_c}$ and

^a e-mail: suhoun@phya.yonsei.ac.kr

^b e-mail: yasui@post.kek.jp

Table 1. Baryon mass relations. The first column is fit to experiments.

Mass	$M_\Delta - M_N$	$M_\Sigma - M_\Lambda$	$M_{\Sigma_c} - M_{\Lambda_c}$	$M_{\Sigma_b} - M_{\Lambda_b}$
Form.	$\frac{3C_B}{2m_u^2}$	$\frac{C_B}{m_u^2}(1 - \frac{m_u}{m_s})$	$\frac{C_B}{m_u^2}(1 - \frac{m_u}{m_c})$	$\frac{C_B}{m_u^2}(1 - \frac{m_u}{m_b})$
Fit	290 MeV	77 MeV	154 MeV	180 MeV
Exp.	290 MeV	75 MeV	170 MeV	192 MeV

$M_{\Sigma_b} - M_{\Lambda_b}$ are well reproduced. This is in no way an attempt to make the best fit, but the point is that with typically accepted constituent quark masses, the mass splittings that are sensitive to the color-spin interaction are well reproduced with a single parameter. As can be seen from Table 2, the fit is equally good for the meson masses with $C_M/m_u^2 = 635$ MeV. Here, even the $B^* - B$ mass difference fits well within the scheme. From these fits, we find that C_M is about 3 times larger than C_B , reflecting that the quark and anti-quark correlation is about 3 times stronger than that between two quarks. The reason why the wave function effect can be treated as a constant is a dynamical question, and some insight can be obtained by looking at various potentials [20]. Within a naive bag model description, the wave function at the origin is only determined by the size of the Bag. Hence, assuming a constant bag radius one finds a constant strength of the wave function at the origin.

Table 2. Meson mass relations. The first column is fit to experiments.

Mass	$M_\rho - M_\pi$	$M_{K^*} - M_K$	$M_{D^*} - M_D$	$M_{B^*} - M_B$
Form.	$\frac{C_M}{m_u^2}$	$\frac{C_M}{m_u m_s}$	$\frac{C_M}{m_u m_c}$	$\frac{C_M}{m_u m_b}$
Fit	635 MeV	381 MeV	127 MeV	41 MeV
Exp.	635 MeV	397 MeV	137 MeV	46 MeV

When both quarks are heavy, the value of C_H is expected to become larger as the strength of the relative wave function at the origin is substantially increased. Fitting instead its value to the mass difference between J/ψ and η_c , we find $C_{c\bar{c}}/m_c^2 = 117$ MeV. Assuming that the corresponding attraction between charmed diquark is three times smaller than that between the charm quark-antiquark pair as in the case of light quarks, we have $C_{cc}/m_c^2 = 39$ MeV. In the mass difference between Υ and η_b , in which the latter has been recently observed by BaBar [21], we also find $C_{b\bar{b}}/m_b^2 = 71.4$ MeV, and in turn $C_{bb}/m_b^2 = 23.8$ MeV. Therefore, we find C_{bb} is much larger than C_{cc} . Concerning the value of C_{bc} , there is no available experimental data for the mass difference between B_c^* and B_c . However, we also expect that C_{cb} has flavor dependence and given by an value interpolating between that for the cc and the bb diquarks. In any case, we find the flavor dependence induces the larger value of C_H for heavier flavors. Therefore, we could introduce additional mass dependence in C_B and in C_M by fitting the mass differences in the strange, charm and bottom hadrons from Table I and II, respectively. However, these introduce only minor changes

in the analysis to follow, and therefore we will just use the mass independent C_H 's obtained above.

3 Stable multi-quark configuration

Using above parameters, one notes that the scalar diquark has a large attractive contribution to the mass. Any multi-quark configuration that are stable should be the ones that are composed of such configurations.

Table 3. Type of attractive scalar diquarks ($q_1 q_2$) composed of quark q_1 and q_2 . The second line is their binding in MeV due to the color-spin force $-\frac{3}{4} \frac{C_B}{m_{q_1} m_{q_2}}$.

(ud)			
-144.75			
(us)	(ds)		
-86.85	-86.85		
(uc)	(dc)	(sc)	
-28.95	-28.95	-17.37	
(ub)	(db)	(sb)	(cb)
-9.23	-9.23	-5.54	-1.84

Table 3 lists the possible diquarks and their attractions due to Eq. (1) with parameters obtained above. Candidates for stable multi-quark configurations can be obtained by combining diquarks and possible color source to make them color singlet.

3.1 Heavy tetraquark

Tetraquarks are obtained by combining a diquark and an antidiquark in Table 3. Let us start by looking at the tetraquark with total spin zero containing the (ud) diquark and an anti-diquark of $(\bar{q}_1 \bar{q}_2)$; the configuration will be called $T_{q_1 q_2}^0$. In the following discussions, we will only consider quarks in the s -state, and assume that the color-spin effect is as given in Eq. (1) with constant coefficients determined above. To investigate the stability of the $T_{q_1 q_2}^0$ against strong decay, one has to consider its mass difference with the two pseudo scalar mesons $M_1(u\bar{q}_1)$ and $M_2(d\bar{q}_2)$, which we call the binding energy $B_{T_{q_1 q_2}^0}$,

$$\begin{aligned}
 B_{T_{q_1 q_2}^0} &= m_{T_{q_1 q_2}^0} - m_{M_1} - m_{M_2} \\
 &= -\frac{3}{4} C_B \left(\frac{1}{m_u^2} + \frac{1}{m_{q_1} m_{q_2}} \right) \\
 &\quad + \frac{3}{4} C_M \left(\frac{1}{m_u m_{q_1}} + \frac{1}{m_u m_{q_2}} \right). \quad (3)
 \end{aligned}$$

Since phenomenologically C_M is about 3 times larger than C_B , the combinations of diquarks which make Eq. (3) negative are very limited. Only when $q_1 = c$ and $q_2 = b$, the binding energy becomes $B_{T_{cb}^0} = -21.25$ MeV, and the configuration $(ud\bar{c}\bar{b})$ which we call T_{cb}^0 will be stable against

strong decay. We note that more realistically the binding energy of a diquark with two heavy quarks could be larger as noted previously. Therefore the binding energy we have obtained will be the lower bound.

This result can be understood simply in the heavy quark limit. The binding energy of T_{cb}^0 is supplied mainly from the binding energy of (ud) diquark, because the binding energy of (cb) diquark is suppressed by $1/m_c m_b$. On the other hand, the binding energies of the two meson states, $u\bar{c}$ and $d\bar{b}$, are also suppressed by $1/m_c$ and $1/m_b$, respectively. Therefore, the (ud) diquark is regarded as an attractive force which stabilizes T_{cb}^0 . Then, we may ask if T_{cb}^0 including (us) or (ds) diquarks can be stable, because s quark may be considered to belong to the light quarks. We find that the configuration $us\bar{c}\bar{b}$ could be stable against $B^+ + D_s^-$ ($B_{T_{cb}^0(us)}^0 = -1.1$ MeV), however it is unstable against $\bar{D}^0 + B_s^0$ ($B_{T_{cb}^0(us)}^0 = +24.8$ MeV) because of the relatively weaker binding in the (us) diquark. Therefore, T_{cb}^0 will not form $\mathbf{3}_f$ multiplet in light flavor $SU(3)_f$ symmetry.

Another tetraquark that could be stable is T_{cc}^1 that was discussed before within the present model in Ref. [18]. T_{cc}^1 is a state with scalar diquark (ud) and axial vector diquark (cc) , hence the total quantum number is $J^P = 1^+$. This state can not decay strongly to two pseudo scalar mesons, but can to a pseudo scalar meson and a vector meson. Therefore, although the charmed anti-diquark is repulsive, the attractive (ud) diquark gives more binding than the pseudo scalar and vector mesons. That is,

$$\begin{aligned} B_{T_{cc}^1} &= m_{T_{cc}^1} - m_D - m_{D^*} \\ &= -\frac{3}{4} \frac{C_B}{m_u^2} + \frac{1}{4} \frac{C_B}{m_c^2} + \frac{3}{4} C_M \left(\frac{1}{m_u m_c} - \frac{1}{3m_u m_c} \right) \\ &\sim -79.3 \text{ MeV}. \end{aligned} \quad (4)$$

In the same way, T_{bb}^1 can also be a stable tetraquark, whose binding energy is given by

$$\begin{aligned} B_{T_{bb}^1} &= m_{T_{bb}^1} - m_B - m_{B^*} \\ &= -\frac{3}{4} \frac{C_B}{m_u^2} + \frac{1}{4} \frac{C_B}{m_b^2} + \frac{3}{4} C_M \left(\frac{1}{m_u m_b} - \frac{1}{3m_u m_b} \right) \\ &\sim -124.3 \text{ MeV}. \end{aligned} \quad (5)$$

We note that T_{cc}^1 containing (us) or (ds) diquark instead of (ud) diquark can also be stable as shown in Table 4. For example, $T_{cc}^1(us)$ has binding energy -8.7 MeV against $\bar{D}^0 + D_s^{*-}$, and $T_{bb}^1(us)$ has -62.3 MeV against $B^+ + B_s^{*0}$. Therefore, $T_{cc}^1(bb)$ will form $\mathbf{3}_f$ multiplet.

In between T_{cc}^1 and T_{bb}^1 , T_{cb}^1 with spin triplet (cb) diquark is also interesting. The binding energy is obtained similarly as shown in Table 4. In this case, T_{cb}^1 with (ud) diquark is stable, while T_{cb}^1 with (us) and (ds) diquarks are not. Therefore, T_{cb}^1 would not form $\mathbf{3}_f$ multiplet, as only T_{cb}^1 is bound. T_{cb}^1 is regarded as an excited state of T_{cb}^0 , whose mass splitting is given by $C_B/m_c m_b$. Therefore, if both T_{cb}^0 and T_{cb}^1 are observed and their mass splitting fits well to $C_B/m_c m_b$, then it confirms that T_{cb} is really a

Table 4. The binding energy $B_{T_{cc(bb,cb)}^1} = m_{T_{cc(bb,cb)}^1} - m_M - m_{M^*}$ of $T_{cc(bb,cb)}^1$ against decay to pseudoscalar and vector mesons, M and M^* , as shown below. The unit is in MeV. The tetraquarks can be stable for $B_{T_{cc(bb,cb)}^1} < 0$.

T_{cc}^1	$ud\bar{c}\bar{c}$	$us\bar{c}\bar{c}$	$ds\bar{c}\bar{c}$
	-79.3 $\bar{D}^0 + D^{*-}, \bar{D}^{*0} + D^-$	-8.7 $\bar{D}^0 + D_s^{*-}$	-8.7 $D^- + D_s^{*-}$
T_{bb}^1	$udbb$	$usbb$	$dsbb$
	-124.3 $B^+ + B^{*0}, B^{*+} + B^0$	-62.3 $B^+ + B_s^{*0}$	-62.3 $B^0 + B_s^{*0}$
T_{cb}^1	$ud\bar{c}\bar{b}$	$us\bar{c}\bar{b}$	$ds\bar{c}\bar{b}$
	-59.0 $B^{*+} + D^-, B^{*0} + \bar{D}^0$	+2.9 $B_s^{*0} + D^0$	+2.9 $B_s^{*0} + D^-$

tetraquark, and not a bound molecular state, which could otherwise be another possibility.

The presence of T_{cb}^0 and T_{cc}^1 can be observed experimentally through their hadronic weak decay [18]. T_{cb}^0 contains the components of $\bar{D}^0 + B^0$ or $D^- + B^+$, whose available decay modes are listed in Ref. [26]. For example, through the weak decay $\bar{D}^0 \rightarrow K^+\pi^-$ and $B^0 \rightarrow K^+\pi^-$, we could observe $T_{cb}^0 \rightarrow K^+\pi^- + K^+\pi^-$, as far as T_{cb}^0 is really stable against the strong decay. While the expected branching ratio for the particular hadronic decay mode is very small because of the small branching ratio of $B^0 \rightarrow K^+\pi^-$, we need to search for such modes to reconstruct the mass of T_{cb}^0 . Similarly, T_{cc}^1 contains $\bar{D}^0 + D^{*-}$ or $\bar{D}^{*0} + D^-$. Then, using $D^* \rightarrow D\pi$, an interesting decay mode is given as $T_{cc}^1 \rightarrow K^+\pi^- + K^+\pi^- + \pi^-$ [18]. In the same way, T_{bb}^1 and T_{cb}^1 contain $B + B^*$ and $D + B^*$, respectively. However, the prediction of decay modes are not easy at present because the available decay modes of B^* are very limited.

3.2 Heavy pentaquark

Pentaquarks are obtained by taking any two diquarks as given in Table 3 together with an antiquark. Stability in this case is guaranteed when the binding energy of the pentaquark configuration is larger than the largest sum of baryon and meson binding energies. The diquarks should be of different types as they have to be in color antisymmetric configuration so that it combines with the antiquark into a color singlet baryon. For the pentaquark to be stable, it would be the best to start with the diquarks with the largest bindings; that is (ud) and (us) . Combining them with an antiquark q , the binding energy of the pentaquark Θ_{qs} ($udus\bar{q}$) becomes,

$$\begin{aligned} B_{\Theta_{qs}} &= m_{\Theta_{qs}} - m_A - m_{u\bar{q}} \\ &= -\frac{3}{4} \frac{C_B}{m_u^2} - \frac{3}{4} \frac{C_B}{m_u m_s} \\ &\quad + \frac{3}{4} \frac{C_B}{m_u^2} + \frac{3}{4} \frac{C_M}{m_u m_q}. \end{aligned} \quad (6)$$

The first and third term cancels. Since C_M is about larger than C_B by a factor 3, it is clear that we will have binding only if $m_q > \frac{C_M}{C_B} m_s > 3m_s$. Therefore, the binding is slightly positive for $q = c$ with 8.4 MeV as shown in Table 5. For $q = b$, one can certainly expect a bound Θ_{bs} whose binding is -56.4 MeV [18]. The large attraction gained from recombining it into a meson and a baryon is the reason why the light pentaquark can not be stable, as was shown in a more detailed constituent quark model [23].

Next, we consider the configuration of (us) and (ds) diquarks with anti-quark $\bar{q}$, which we denote as $\Theta_{qss}(usds\bar{q})$. In this case we have two possible final states; $\Xi + u\bar{q}$ and $\Lambda + s\bar{q}$. The binding energy for the former state is given by

$$\begin{aligned} B_{\Theta_{qss}} &= m_{\Theta_{qss}} - m_{\Xi} - m_{u\bar{q}} \\ &= -2 \times \frac{3}{4} \frac{C_B}{m_u m_s} + \frac{3}{4} \frac{C_B}{m_u m_s} + \frac{3}{4} \frac{C_M}{m_u m_q}, \end{aligned} \quad (7)$$

and that for the latter state by

$$\begin{aligned} B_{\Theta_{qss}} &= m_{\Theta_{qss}} - m_{\Lambda} - m_{s\bar{q}} \\ &= -2 \times \frac{3}{4} \frac{C_B}{m_u m_s} + \frac{3}{4} \frac{C_B}{m_u^2} + \frac{3}{4} \frac{C_M}{m_s m_q}. \end{aligned} \quad (8)$$

The numerical result is shown in Table 5. In these configurations, we again find that the pentaquark becomes more stable as the included antiquark becomes heavier. Therefore, Θ_{cs} and Θ_{css} are slightly unstable, while Θ_{bs} and Θ_{bss} are stable. Nonetheless, it is worth searching for all these states in the experiments.

The flavor structure of the stable pentaquark is obtained by following light flavor $SU(3)_f$ symmetry. In the present discussion, the stable pentaquark contains (ud) , (us) and (ds) diquarks together with a heavy quark, say $q = b$. Because these diquarks belong to $\bar{\mathbf{3}}_f$ representation in $SU(3)_f$, the flavor decomposition is obtained as

$$\bar{\mathbf{3}}_f \times \bar{\mathbf{3}}_f = \mathbf{3}_f^a + \bar{\mathbf{6}}_f^s.$$

In our diquark model, $\bar{\mathbf{6}}_f^s$ vanishes because it is symmetric under the exchange of flavors and hence can be canceled by anti-symmetry in color. Therefore, the pentaquarks obtained above belong to the $\mathbf{3}_f^a$ representation.

Table 5. The binding energy of pentaquarks $B_{\Theta_{qs}} = m_{\Theta_{qs}} - m_M - m_B$ ($B_{\Theta_{qs}} = m_{\Theta_{qs}} - m_M - m_B$) of $\Theta_{qs}(udus\bar{q})$ ($\Theta_{qs}(usds\bar{q})$) for $q = u, s, c$ and b , respectively, against decays to meson M and baryon B states. The unit is in MeV. The pentaquarks can be stable for $B_{\Theta_{qs}}(\Theta_{qss}) < 0$.

Θ	$M+B$	$\bar{u}$	$\bar{s}$	$\bar{c}$	b
$udus\bar{q}$	$uds+u\bar{q}$	389.4	198.9	8.4	-56.4
$usds\bar{q}$	$ds s+u\bar{q}$	389.4	198.9	8.4	-56.4
	$uds+s\bar{q}$	256.8	142.5	28.4	-10.7

The pentaquarks have easily identifiable decay modes. For example, $\Theta_{cs}(udds\bar{c}) \rightarrow \Lambda + K^+\pi^-$ [18], and Θ_{bs}

$(udds\bar{b}) \rightarrow \Lambda + K^+\pi^-$, as modes with the least number of particles in the final states. Similarly, $\Theta_{css} \rightarrow \Lambda + \pi^+\pi^-\pi^-$, and $\Theta_{bss} \rightarrow \Lambda + \pi^+\pi^-\pi^- + \pi^+$.

3.3 Heavy dibaryon

Dibaryons are obtained by taking three diquarks and combining them into a color singlet combination. Considering the color anti-triplet scalar diquarks $(q_i q_j)$ in Table 3, the wave function is given as,

$$\epsilon_{abc}(q_1 q_2)^a (q_3 q_4)^b (q_5 q_6)^c, \quad (9)$$

where a, b, c are the color indexes. The total number of states which can be obtained from these wave functions are 120. However, only a few of them are stable against strong decay to two baryons. To understand which configurations are stable, let us consider the decay of the dibaryon into two baryons in the following way. First of all, let us assume that the diquark $(q_1 q_2)$ is the most stable one among the three diquarks in Eq. (9). If there are two most stable diquarks, one of them is assigned to $(q_1 q_2)$. When the dibaryon decays to two baryons, there should be a process where some diquarks in the dibaryon are broken into quarks. In the present discussion, because $(q_1 q_2)$ is the most stable diquark, the remaining two diquarks, $(q_3 q_4)$ and $(q_5 q_6)$, can be broken. First possible process occurs when one diquark is broken. When $(q_5 q_6)$ is more loosely bound than $(q_3 q_4)$, then the former diquark is broken to a pair of single quarks q_5 and q_6 . Then they are combined with $(q_1 q_2)$ and $(q_3 q_4)$ to form two baryons that can be represented as either $(q_1 q_2)q_5$ and $(q_3 q_4)q_6$, or $(q_1 q_2)q_6$ and $(q_3 q_4)q_5$. This process is shown by the following equation,

$$\begin{aligned} \text{conf}_1 &= (q_1 q_2)q_5 + (q_3 q_4)q_6, \\ &\text{or } (q_1 q_2)q_6 + (q_3 q_4)q_5. \end{aligned} \quad (10)$$

Second possible process occurs when two diquarks are broken. In this case, the diquarks $(q_3 q_4)$ and $(q_5 q_6)$ are first broken into q_3, q_4, q_5 and q_6 . Second, when the most stable diquark among possible combinations of these four quarks is $(q_3 q_5)$, then they are combined to two possible final baryon states that are either $(q_1 q_2)q_4$ and $(q_3 q_5)q_6$, or $(q_1 q_2)q_6$ and $(q_3 q_5)q_4$. This process is shown by,

$$\begin{aligned} \text{conf}_2 &= (q_1 q_2)q_4 + (q_3 q_5)q_6, \\ &\text{or } (q_1 q_2)q_6 + (q_3 q_5)q_4. \end{aligned} \quad (11)$$

The stability of dibaryons is determined by the mass difference between the dibaryon and two baryon states. Namely, when the mass of dibaryon is smaller than those of conf_1 and conf_2 , then such dibaryon can be stable against the strong decay. Now we investigate the stability of the configurations of conf_1 or conf_2 .

Let us first consider the stability of dibaryon against the configuration conf_1 . In the present diquark model, the diquark $(q_5 q_6)$ is a bound state for any flavor combinations with finite binding energy as given in Table 3. Therefore, the mass of dibaryon is always smaller than that of conf_1 by the binding energy of $(q_5 q_6)$. Consequently, we obtain the following rule.

- i) Any dibaryon constructed from Table 3 is stable against the decay to conf_1 .

However, the dibaryons are not necessarily stable against the decay to conf_2 . To see this, let us analyze the stability against the decay to conf_2 in Eq. (11). For this purpose, we consider the difference between the binding energies of the dibaryon and conf_2 . Omitting the common binding energy supplied from (q_1q_2) , the binding energy of the dibaryon is given by

$$\text{B.E.} \left[(q_3q_4) + (q_5q_6) \right] = -\frac{3}{4} \frac{C_B}{m_3m_5} \left(\frac{m_5}{m_4} + \frac{m_3}{m_6} \right), \quad (12)$$

while that of conf_2 is given by

$$\text{B.E.} \left[(q_3q_5) \right] = -\frac{3}{4} \frac{C_B}{m_3m_5}. \quad (13)$$

As can be seen above, the dibaryon is stable if the following condition is satisfied,

$$\frac{m_5}{m_4} + \frac{m_3}{m_6} > 1. \quad (14)$$

From this equation, we obtain two rules to realize a stable dibaryon.

- ii-a) First, we considering the case where the two diquarks are in the same row in Table 3; $q_4 = q_6$. Then condition (14) reduces to

$$m_3 + m_5 > m_4. \quad (15)$$

Because two diquarks are taken from a single row in Table 3, we have the following possibilities; $\{(q_3q_4), (q_5q_6)\} = \{(us), (ds)\}$ for $q_4 = s$, $\{(uc), (dc)\}$, $\{(uc), (sc)\}$ and $\{(dc), (sc)\}$ for $q_4 = c$, $\{(ub), (db)\}$, $\{(ub), (sb)\}$, $\{(ub), (cb)\}$, $\{(db), (sb)\}$, $\{(db), (cb)\}$ and $\{(sb), (cb)\}$ for $q_4 = b$.

Note $q_3 \neq q_5$ from Eq. (11). Among them, condition (15) is satisfied only for $\{(q_3q_4), (q_5q_6)\} = \{(us), (ds)\}$. In other words, diquarks which contain at least two diquarks from either the third or the fourth row of Table 3, such as $\{uc, dc\}$, can not be stable against conf_2 .

- ii-b) Second, we consider the case where the two diquarks are in different row of Table 3; $q_4 \neq q_6$. In this case however, one finds that no configuration satisfies Eq. (14) when q_3 and q_5 are different light quarks.

Combining these two rules, one can obtain stable dibaryon configurations by starting from either the first or second row and picking up the remaining two diquarks by moving down in rows either straight along a single column or to different columns with Eq. (14) satisfied. Following these rules about decays to conf_1 and conf_2 , it is easy to find that there are 16 stable dibaryons containing (ud) diquark and 9 more containing (us) or (ds) diquarks. The only exception to this rule is the $(ud)(sc)(sb)$ as it can decay into $(us)c + (ds)b$ type baryons. Table 6 lists all the possible configurations and the stable dibaryons whose binding energy are negative.

In Table 6, the stable dibaryons are categorized according to their contents of diquarks as represented in the

first column; $(qq)(qq)(qq)$, $(qq)(qq)(qQ)$, $(qq)(qQ)(qQ')$, $(qq)(qq)(QQ')$ and $(qq)(qQ)(QQ')$ with $q = u, d, s$ and $Q, Q' = c, b$ which may be called H , H_Q , $H_{Q'Q}$, $H'_{Q'Q}$ and $H_{Q'QQ}$, respectively. The second column indicates the flavor content through five integers $N_u N_d N_s N_c N_b$ representing the number of quark with u, d, s, c and b flavors respectively. The third column is the configurations of diquarks in dibaryons. Here, for short description, we use qq instead of (qq) . The fourth, fifth and sixth columns are isospin, its z component and strangeness, respectively. The seventh column indicates the configurations of the two baryon states with lowest threshold. The binding energy of the dibaryon against the two baryon threshold is shown as their mass difference in the last column.

The $(qq)(qq)(qq)$ type becomes stable from the rule ii-a) for conf_2 , while the other configurations are stable from the rule i) for conf_1 . We mention that several dibaryons which are not accounted from the rules i), ii-a) and ii-b) are also stable as components in isospin multiplets as shown in Table 6. They are the flavor $N_u N_d N_s N_c N_b = 22110$ with $I = 1$ and $I_z = 0$ in $(qq)(qq)(qQ)$ ($Q = c$), 22101 with $I = 1$ and $I_z = 0$ in $(qq)(qq)(qQ)$ ($Q = b$), 21111 and 12111 with $I = 3/2$ and $I_z = \pm 1/2$ in $(qq)(qQ)(qQ)$, and 11121 with $I = 1$, $I_z = 0$ in $(qq)(qQ)(QQ')$.

Among these dibaryons, the following two dibaryon configurations are the most deeply bound states and are worth mentioning in details. The first one is the H dibaryon [4] with diquark configurations of $(ud)(us)(ds)$. Again, the binding energy of the H dibaryon can be obtained by comparing its color-spin interaction to that of two Λ 's,

$$\begin{aligned} B_H &= m_H - m_\Lambda - m_\Lambda \\ &= -\frac{3}{4} \frac{C_B}{m_u^2} - 2 \frac{3}{4} \frac{C_B}{m_u m_s} \\ &\quad + \frac{3}{4} \frac{C_B}{m_u^2} + \frac{3}{4} \frac{C_B}{m_u^2}. \end{aligned} \quad (16)$$

The first term cancels the third term. Therefore, the stability of H dibaryon depends on whether it is energetically favorable to have $(us) + (ds)$ diquarks for a strong (ud) diquark, which in fact is the rule ii-a) above. Substituting the constants in our model, we find a binding of $B_H = -29$ MeV, which is consistent with most of the model calculations [24].

The second one is the dibaryon of the $(qq)(qq)(qQ)$ type with $Q = c$ containing a single charm, which we will call the H_c dibaryon. The most attractive dibaryon in such configuration is $(ud)(us)(uc)$, $(ud)((us)(dc) + (ds)(uc))$ and $(ud)(ds)(dc)$ in the isospin $I = 1$ multiplet. Concerning $(ud)(us)(uc)$, the binding energy of H_c can be investigated by comparing its color-spin interaction to that of the proton (udu) and Ξ_c (usc),

$$\begin{aligned} B_{H_c} &= m_{H_c} - m_p - m_{\Xi_c} \\ &= -\frac{3}{4} \frac{C_B}{m_u^2} - \frac{3}{4} \frac{C_B}{m_u m_s} - \frac{3}{4} \frac{C_B}{m_u m_c} \\ &\quad + \frac{3}{4} \frac{C_B}{m_u^2} + \frac{3}{4} \frac{C_B}{m_u m_s} \\ &= -\frac{3}{4} \frac{C_B}{m_u m_c}. \end{aligned} \quad (17)$$

The same binding energy is also obtained for the other isospin partners. In another view, it can be regarded that the binding energy of these H_c dibaryons is supplied from the binding energy of (uc) or (dc) diquarks. As a result, one obtains $B_{H_c} = -29$ MeV and a stable H_c dibaryon. This is a new prediction and could be searched in future heavy ion collisions, where large number of charm quarks are expected to be produced and weakly decaying particle identifiable through vertex detectors [18]. As further possible bound states, $(ud)(us)(sc)$ and $(ud)(ds)(sc)$ in $I = 1/2$ and $(us)(ds)(sc)$ in $I = 0$ all have binding energies of -17 MeV. This binding energy is supplied from the binding of the (sc) diquark, and hence is smaller than that of H_c dibaryon. Nevertheless, these configurations are additional candidates for stable dibaryons containing a single charm.

The dibaryons stated above have relatively large binding energies as the H dibaryon. The diquark model also predicts other possible dibaryon states which have smaller binding energies. In the $(qq)(qq)(qQ)$ type, we obtain stable configurations for $Q = b$ (H_b) with binding energies equal to -9 MeV for $I = 1$, and -5 MeV for $I = 1/2$ and for $I = 0$, as shown in Table 6. The binding energy is reduced in comparison with $Q = c$, simply because the (ub) , (db) and (sb) diquarks are more loosely bound. As for the dibaryons containing two heavy quarks, the $(qq)(qq)(qQ')$ and $(qq)(qq)(QQ')$ types with $Q = c$ and $Q' = b$ (H_{bc} and H'_{bc}) can also be stable. For the former, the binding energies are obtained as -9 MeV for $I = 1$ and $I = 3/2$, and -5 MeV for $I = 1/2$ which bindings are supplied from each of (ub) , (db) and (sb) diquarks. For the latter, the binding energies are -1.8 MeV for $I = 1/2$ and $I = 0$ which are supplied from the (cb) diquark. The binding energy of (cb) diquark is very small in comparison with other diquarks containing light flavors. Therefore, it supplies only small binding energy for the dibaryons. Lastly, we can have dibaryons containing three heavy quarks as $(qq)(qQ)(QQ')$ with $Q = c$ and $Q' = b$ (H_{bcc}). However, their binding energies are also small because they are supplied from the (cb) diquark.

According to the $SU(3)_f$ light flavor symmetry, the dibaryons in Table 6 belong to the following multiplets. Noting (qq) diquark belongs to $\bar{\mathbf{3}}_f$, the light flavor multiplets of the $(qq)(qq)(qq)$ type are obtained by $\bar{\mathbf{3}}_f \times \bar{\mathbf{3}}_f \times \bar{\mathbf{3}}_f = \mathbf{1}_f + \mathbf{8}_f + \mathbf{8}_f + \mathbf{10}_f$. Then the H dibaryon $(ud)(us)(ds)$ belongs to $\mathbf{1}_f$.

For $(qq)(qq)(qQ)$ with $Q = c$ or b (H_c or H_b), noting (qQ) diquark belongs to $\mathbf{3}_f$, the light flavor is decomposed as

$$\bar{\mathbf{3}}_f \times \bar{\mathbf{3}}_f \times \mathbf{3}_f = (\bar{\mathbf{3}}_f^a + \bar{\mathbf{6}}_f^s) \times \mathbf{3}_f = \bar{\mathbf{3}}_f^a + \mathbf{6}_f + \bar{\mathbf{3}}_f^s + \mathbf{15}_f.$$

Here a (s) means anti-symmetry (symmetry) for a pair of two diquarks (qq) 's in $(qq)(qq)(qQ)$. Because of the anti-symmetry of color in the wave function in Eq. (9), $\bar{\mathbf{3}}_f^a$ and $\mathbf{15}_f$ which contain identical pair of diquarks in symmetric flavor $\bar{\mathbf{6}}_f^s$ vanish, and only $\bar{\mathbf{3}}_f^s$ and $\mathbf{6}_f$ which contain anti-symmetric flavor $\mathbf{3}_f^a$ survive. Concerning mixing of $\bar{\mathbf{3}}_f^s$ and $\mathbf{6}_f$, some comments are in order. First, in $S = -1$ sector, the wave function in the $I = 0$ configuration in $\bar{\mathbf{3}}_f^a$ is given

as

$$(ud) \{ (us)(dQ) - (ds)(uQ) \},$$

while that of $I_z = 0$ in $I = 1$ in $\mathbf{6}_f$, which weight vector is on the same place, is given as

$$(ud) \{ (us)(dQ) + (ds)(uQ) \},$$

with appropriate normalization constants. As far as the isospin is conserved exactly, these two states are independent. The latter dibaryon with $I = 1$ is stable against the decay to $\Lambda + \Lambda_Q$ because isospin symmetry forbids this decay. However, the former dibaryon with $I = 0$ is not stable, but decays to $\Lambda + \Lambda_Q$. Therefore, in $S = -1$ sector, we have stable dibaryons in $I = 1$. Second, in $S = -2$ sector, the $I = 1/2$ configuration in $\bar{\mathbf{3}}_f^a$ are generally mixed with those in $\mathbf{6}_f$. For example, the $I_z = 1/2$ components are given by

$$(ud)(us)(sQ) \mp (us)(ds)(uQ)$$

for $\bar{\mathbf{3}}_f^a$ and $\mathbf{6}_f$, respectively. However, from the rule to obtain the stable dibaryon, the dibaryon $(us)(ds)(uQ)$ is not stable against $\Lambda + \Xi_Q$ and $\Lambda_Q + \Xi$. Therefore, by considering a superposition of $\bar{\mathbf{3}}_f^a$ and $\mathbf{6}_f$ and eliminating $(us)(ds)(uQ)$, we obtain the stable dibaryon $(ud)(us)(sQ)$. Third, in $S = -3$ sector, the (us) and (ds) pair in $(us)(ds)(sQ)$ is allowed to be $I = 0$ from anti-symmetry of color configuration. The resulting weight diagrams are shown in Fig. 1 and 2 for $Q = c$ and b , respectively.

Similarly, other multiplets for the configurations with two or three heavy quarks are obtained. As for $(qq)(qQ)(qQ')$ with $Q = c$ and $Q' = b$ (H_{bc}), the light flavor is decomposed by

$$\bar{\mathbf{3}}_f \times \mathbf{3}_f \times \mathbf{3}_f = \bar{\mathbf{3}}_f \times (\bar{\mathbf{3}}_f^a + \mathbf{6}_f^s) = \mathbf{3}_f^a + \bar{\mathbf{6}}_f + \mathbf{3}_f^s + \mathbf{15}_f.$$

We note that $\mathbf{3}_f^s$ and $\mathbf{15}_f$ containing symmetric flavor $\mathbf{6}_f^s$ can survive, because (qQ) cannot be identical to (qQ') due to $Q \neq Q'$. From the configurations in Table 6, we find that the stable dibaryons contain pairs of (qQ) and (qQ') with identical q , which are assigned to $\mathbf{6}_f^s$. Hence, these dibaryons belong to $\mathbf{3}_f^s$ or $\mathbf{15}_f$. Here, we show that only $\mathbf{15}_f$ is relevant to the stable dibaryons. First, in $S = -1$ sector, the $I = 1/2$ configuration in $\mathbf{3}_f^s$ is at the same position in the weight diagram with the $I_z = \pm 1/2$ components in the $I = 3/2$ configuration in $\mathbf{15}_f$. However, the $I = 1/2$ configuration is not stable, because, for example, the $I_z = 1/2$ component in $I = 1/2$ can decay to $\Lambda_b + \Xi_c^+$ (or $\Lambda_c + \Xi_b^0$). Hence the $S = -1$ sector in $\mathbf{3}_f^s$ is irrelevant. Second, in $S = -2$ sector, any components in $I = 0$ in $\mathbf{3}_f^s$ and in $I = 1$ in $\mathbf{15}_f$ are not stable. In fact we have already discussed that $(ud)(sc)(sb)$ configuration is not stable. Therefore, $\mathbf{3}_f^s$ can be discarded, and so can the $I = 1$ configuration in $S = -2$ in $\mathbf{15}_f$. Concerning the $S = 0$ sector, we note that the $I_z = 0$ component of $I = 1$ cannot decay to $\Lambda_c + \Lambda_b$ because of isospin conservation, and hence can be stable. The resulting weight diagram is shown in Fig. 3. Here, the black blobs indicate the stable dibaryons, while the white ones indicate the unstable ones.

As for the $(qq)(qq)(QQ')$ type with $Q = c$ and $Q' = b$ (H'_{bc}), the light flavor is decomposed as

$$\bar{\mathbf{3}}_f \times \bar{\mathbf{3}}_f = \mathbf{3}_f^a + \bar{\mathbf{6}}_f^s.$$

Because a pair of two (qq) 's in $\bar{\mathbf{6}}_f^s$ vanishes due to anti-symmetry in color, only $\mathbf{3}_f^a$ can survive because of isospin conservation. The weight diagram of $(qq)(qq)(QQ')$ is shown in Fig. 4.

Lastly, for dibaryons with three heavy quarks such as $(qq)(qQ)(QQ')$ type with $Q = c$ and $Q' = b$ (H_{bcc}), the light flavor is decomposed as

$$\bar{\mathbf{3}}_f \times \mathbf{3}_f = \mathbf{1}_f + \mathbf{8}_f.$$

In $S = -1$ sector, the $I = 0$ configurations in $\mathbf{1}_f$ are superposition of $((us)(dQ) - (ds)(uQ))(QQ')$ and $(ud)(sQ)(QQ')$, which are constructed from (us) , (ds) and (ud) diquarks. Among them, however, the dibaryons containing (us) or (ds) are not stable against $\Lambda_b + \Omega_{cc}^+$ (or $\Lambda_c + \Omega_{cb}^0$), and hence $(ud)(sQ)(QQ')$ containing only (ud) diquark is stable. On the other hand, the $I = 1$ configuration of $((us)(dQ) + (ds)(uQ))(QQ')$ in $\mathbf{8}_f$ can survive. In $S = -2$, there are no stable dibaryons. As a result, the weight diagram is obtained in Fig. 5.

Now, let us discuss possible decay modes from these dibaryons for experiments. Concerning H dibaryon, the observation has been studied by weak decay, for example, by $H \rightarrow \Lambda + \Lambda \rightarrow p\pi^- + p\pi^-$. Similarly, we investigate possible decay modes of the stable dibaryon including heavy quarks.

The $(qq)(qq)(qQ)$ with $Q = c$, which would be most stable state, can have the decay modes as summarized in Table 7. For example, the wave function of the $(ud)(us)(uc)$ with $I_z = 1$ in $I = 1$ would have an overlap with $\Lambda_c^+ + \Sigma^+$ and $p + \Xi_c^+$ with invariant mass 3475 MeV and 3406 MeV, respectively. Then, each component will decay by weak interaction; $\Lambda_c^+ + \Sigma^+ \rightarrow pK^- \pi^+ + p\pi^0$, and $p + \Xi_c^+ \rightarrow p + \Lambda K^- \pi^+ \pi^+$, $\Sigma^+ K^- \pi^+$, $\Xi^0 \pi^+$, $\Xi^- \pi^+ \pi^+$, $\Xi^0 \pi^+ \pi^0$, $\Xi^0 \pi^+ \pi^+$ [26]. We note that, concerning Λ_c , we indicate only $\Lambda_c \rightarrow pK^- \pi^+$ as a simplest decay pattern. The $I_z = -1$ component in $I = 1$ has also an overlap with $\Lambda_c^+ + \Sigma^-$ and $n + \Xi_c^0$ with invariant mass 3483 MeV and 3410 MeV, respectively, and the decay modes of each component are $\Lambda_c^+ + \Sigma^- \rightarrow pK^- \pi^+ + n\pi^-$, and $n + \Xi_c^0 \rightarrow n + pK^- K^- \pi^+$, $\Lambda K^- \pi^+ \pi^+ \pi^-$, $\Xi^- \pi^+$, $\Xi^- \pi^+ \pi^+ \pi^-$, $\Omega^- K^+$. The $I_z = 0$ component in $I = 1$ would contain $\Lambda + \Sigma_c^+$ or $\Lambda_c^+ + \Sigma^-$ because of isospin symmetry, and hence its decay modes are $\Lambda + \Sigma_c^+ \rightarrow p\pi^- + \Lambda_c \pi^0 \rightarrow p\pi^- + pK^- \pi^+ \pi^0$, and $\Lambda_c^+ + \Sigma^- \rightarrow \Lambda \gamma + pK^- \pi^+$. However, in reality, explicit isospin breaking effect would induce strong decay much larger than weak decay. Hence these states would first decay to $H_c \rightarrow \Lambda + \Lambda_c^+$ by explicit isospin breaking, and second they would decay to $p\pi^- + pK^- \pi^+ \pi^0$ by weak interaction. As for $I = 1/2$ and $I = 0$, which would be second most stable states, we obtain the possible decay modes as shown in 7. Note that the isospin $I = 0$ indicates the wave function $\Xi^0 \Xi_c^0 - \Xi^- \Xi_c^+$ and each component is shown in the table.

Similar decay patterns hold for $(qq)(qq)(qQ)$ with $Q = b$ as shown in Table 8. However, because of the lack of exper-




Fig. 1. The weight diagram of $SU(3)_f$ for stable diquark of $qq qq qQ$ ($Q = c$) type (H_c) with strangeness S , isospin I and binding energy B_H . The black blobs indicate the stable dibaryons.




Fig. 2. The weight diagram of $SU(3)_f$ for stable diquark of $qq qq qQ$ ($Q = b$) type (H_b) with strangeness S , isospin I and binding energy B_H . The black blobs indicate the stable dibaryons.

imental data about the decays of several bottom baryons, available decay patterns are limited at present. Nevertheless at least the $I = 1$ and $I = 1/2$ dibaryons can be detectable, while the $I = 0$ dibaryon needs experimental information about the decays of Ξ_b^0 or Ξ_b^- .

As for dibaryons with two or three heavy quarks, only a few candidates are available. In $(qq)(qQ)(qQ')$ with $Q = c$ and $Q' = b$, the $I_z = \pm 1$ components in $I = 1$ and $I = 3/2$ could be investigated by using experimentally known decay patterns as shown in Table 9. In $(qq)(qq)(Q'Q')$ with $Q = c$ and $Q' = b$, the $I_z = \pm 1/2$ components in $I = 1/2$ could be also investigated as shown in Table 10. Lastly, any components in $(qq)(qQ)(Q'Q')$ with $Q = c$ and $Q' = b$ would be difficult because of the lack of experimental information as shown in Table 11.

Thus, the diquark model gives a list of possible H dibaryons, H_c , H_b , H_{bc} , H'_{bc} and H_{bcc} . Among them, the first promising candidate is H_c with $I = 0$, and the second ones are H_c with $I = 1/2$ and 1. The other H dibaryons, H_b , H_{bc} , H'_{bc} and H_{bcc} , are less probable because of their small binding energies. Nevertheless, search of these particles is a challenging subject, not only for theory with more sophisticated analysis by lattice simulation, QCD sum rule, and so on, but also for experiments in high energy facilities.




Fig. 3. The weight diagram of $SU(3)_f$ for stable diquark of $qq qQ qQ'$ ($Q = c$ and $Q' = b$) type (H_{bc}) with strangeness S , isospin I and binding energy B_H . The black blobs indicate the stable dibaryons, while the white blobs do the unstable dibaryons.




Fig. 4. The weight diagram of $SU(3)_f$ for stable diquark of $qq qq QQ'$ ($Q = c$ and $Q' = b$) type (H'_{bc}) with strangeness S , isospin I and binding energy B_H . The black blobs indicate the stable dibaryons.




Fig. 5. The weight diagram of $SU(3)_f$ for stable diquark of $qq qQ QQ'$ ($Q = c$ and $Q' = b$) type (H_{bcc}) with strangeness S , isospin I and binding energy B_H . The black blobs indicate the stable dibaryons, while the white blobs do the unstable dibaryons.

4 Summary

Based on the consideration of the color-spin interaction between diquarks, which describes well the mass splittings between many hadrons and their spin flipped partners, we have systematically investigated possible stable multi-quark configurations. We have shown that the configurations becomes stable only when heavy quarks are included. We have identified several new possibly stable multi-quark

configurations. These are the $T_{cb}^0(ud\bar{c}\bar{b})$ tetraquark, the $\Theta_{bs}(udus\bar{b})$ pentaquark and the $H_c(udusuc)$ dibaryon, and so forth. The representative decay modes for these states are $T_{cb}^0 \rightarrow K^+\pi^- + K^+\pi^-$, $\Theta_{bs} \rightarrow \Lambda K^+\pi^-$ and $H_c \rightarrow \Lambda + \Lambda_c^+ \rightarrow p\pi^- + pK^-\pi^+\pi^0$. If one observes these states, these will be the first explicit flavor multi-quark states, and will be a valuable stepping stone to understanding QCD at larger quark number.

The work of SHL was supported by the Korea Research Foundation KRF-2006-C00011.

References

1. R. L. Jaffe, Phys. Rev. D **15** (1977) 267.
2. K. Abe *et al.* [Belle Collaboration], Phys. Rev. Lett. **100** (2008) 142001 [arXiv:0708.1790 [hep-ex]].
3. R. Mizuk *et al.* [Belle Collaboration], Phys. Rev. D **78** (2008) 072004 [arXiv:0806.4098 [hep-ex]].
4. R. L. Jaffe, Phys. Rev. Lett. **38** (1977) 195 [Erratum-ibid. **38** (1977) 617].
5. J. Schaffner-Bielich and A. P. Vischer, Phys. Rev. D **57** (1998) 4142 [arXiv:nucl-th/9710064].
6. C. Gignoux, B. Silvestre-Brac and J. M. Richard, Phys. Lett. B **193** (1987) 323.
7. H. J. Lipkin, Phys. Lett. B **195** (1987) 484.
8. D. O. Riska and N. N. Scoccola, Phys. Lett. B **299** (1993) 338.
9. Y. s. Oh, B. Y. Park and D. P. Min, Phys. Rev. D **50** (1994) 3350.
10. Y. s. Oh and B. Y. Park, Z. Phys. A **359** (1997) 83.
11. I. W. Stewart, M. E. Wessling and M. B. Wise, Phys. Lett. B **590**, (2004) 185.
12. Y. Sarac, H. Kim and S. H. Lee, Phys. Rev. D **73** (2006) 014009.
13. J. Carlson, L. Heller and J. A. Tjon, Phys. Rev. D **37** (1988) 744.
14. B. Silvestre-Brac and C. Semay, Z. Phys. C **59** (1993) 457.
15. A. V. Manohar and M. B. Wise, Nucl. Phys. B **399** (1993) 17 [arXiv:hep-ph/9212236].
16. R. L. Jaffe and F. Wilczek, Phys. Rev. Lett. **91** (2003) 232003 [arXiv:hep-ph/0307341].
17. M. G. Alford, K. Rajagopal and F. Wilczek, Phys. Lett. B **422** (1998) 247 [arXiv:hep-ph/9711395]; R. Rapp, T. Schafer, E. V. Shuryak and M. Velkovsky, Phys. Rev. Lett. **81** (1998) 53 [arXiv:hep-ph/9711396].
18. S. H. Lee, S. Yasui, W. Liu and C. M. Ko, Eur. Phys. J. C **54** (2008) 259 [arXiv:0707.1747 [hep-ph]].
19. A. De Rujula, H. Georgi and S. L. Glashow, Phys. Rev. D **12** (1975) 147.
20. B. Keren-Zur, Annals Phys. **323** (2008) 631 [arXiv:hep-ph/0703011].
21. B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **101** (2008) 071801 [arXiv:0807.1086 [hep-ex]].
22. J. L. Rosner, Phys. Rev. D **75** (2007) 013009 [arXiv:hep-ph/0611207].
23. E. Hiyama, M. Kamimura, A. Hosaka, H. Toki and M. Yahiro, Phys. Lett. B **633** (2006) 237 [arXiv:hep-ph/0507105].
24. T. Sakai, K. Shimizu and K. Yazaki, Prog. Theor. Phys. Suppl. **137** (2000) 121 [arXiv:nucl-th/9912063].

- 25. M. G. Alford, A. Schmitt, K. Rajagopal and T. Schafer,
Rev. Mod. Phys. **80** (2008) 1455 [arXiv:0709.4635 [hep-ph]].
- 26. C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**
(2008) 1.

Table 6. The mass difference of H -dibaryons and two baryon state B and B' . $B_H = m_H - m_B - m_{B'}$. The H -dibaryons can be stable for $B_H < 0$. $Q \neq Q'$. The unit is in MeV. Contrary to the text, we omit brackets denoting scalar and isoscalar diquark for simplicity.

	flavor	config. of H	I	I_z	S	config. of $B + B'$	B_H [MeV]
$H(qq qq qq)$	22200	$ud us ds$	0	0	-2	$ud s + ud s$	-28.95
$H_c(qq qq qc)$	31110	$ud us uc$	1	1	-1	$ud c + us u, ud u + us c$	-28.95
	22110	$ud \frac{1}{\sqrt{2}}(us dc + ds uc)$	1	0	-1	$ud s + ud c, ud s + ud c$	
	13110	$ud ds dc$	1	-1	-1	$ud c + ds d, ud d + ds c$	
	21210	$ud us sc$	1/2	1/2	-2	$ud c + us s, ud s + us c$	-17.37
	12210	$ud ds sc$	1/2	-1/2	-2	$ud c + ds s, ud s + ds c$	
	11310	$\frac{1}{\sqrt{2}}(us ds - ds us) sc$	0	0	-3	$us s + ds c, us c + ds s$	-17.37
$H_b(qq qq qb)$	31101	$ud us ub$	1	1	-1	$ud u + us b, ud b + us u$	-9.23
	22101	$ud \frac{1}{\sqrt{2}}(us db + ds ub)$	1	0	-1	$ud s + ud b, ud s + ud b$	
	13101	$ud ds db$	1	-1	-1	$ud b + ds d, ud d + ds b$	
	21202	$ud us sb$	1/2	1/2	-2	$ud b + us s, ud s + us b$	-5.54
	12201	$ud ds sb$	1/2	-1/2	-2	$ud b + ds s, ud s + ds b$	
	11301	$\frac{1}{\sqrt{2}}(us ds - ds us) sb$	0	0	-3	$us b + ds s, us s + ds b$	-5.54
$H_{bc}(qq qc qb)$	31011	$ud uc ub$	1	1	0	$ud b + uc u, ud u + uc b$	-9.23
	22011	$ud \frac{1}{\sqrt{2}}(uc db + dc ub)$	1	0	0	$ud b + uc u, ud u + uc b$	
	13011	$ud dc db$	1	-1	0	$ud b + dc d, ud d + dc b$	
	30111	$us uc ub$	3/2	3/2	-1	$us u + uc b, us b + uc u$	-9.23
	21111	$\frac{1}{\sqrt{3}}((us dc + ds uc) ub + us uc db)$	3/2	1/2	-1	$ud b + us c, ud c + us b$	
	12111	$\frac{1}{\sqrt{3}}((us dc + ds uc) db + ds dc ub)$	3/2	-1/2	-1	$ud b + ds c, ud c + ds b$	
	03111	$ds dc db$	3/2	-3/2	-1	$ds b + dc d, ds d + dc b$	
	10311	$us sc sb$	1/2	1/2	-3	$us b + sc s, us s + sc b$	-5.54
	01311	$ds sc sb$	1/2	-1/2	-3	$ds b + sc s, ds s + sc b$	
$H'_{bc}(qq qq bc)$	21111	$ud us cb$	1/2	1/2	-1	$ud b + us c, ud c + us b$	-1.84
	12111	$ud ds cb$	1/2	-1/2	-1	$ud b + ds c, ud c + ds b$	
	11211	$\frac{1}{\sqrt{2}}(us ds - ds us) cb$	0	0	-2	$us b + ds c, us c + ds b$	-1.84
$H_{bcc}(qq qc cb)$	21021	$ud uc cb$	1/2	1/2	0	$ud b + uc c, ud c + uc b$	-1.84
	12021	$ud dc cb$	1/2	-1/2	0	$ud b + dc c, ud c + dc b$	
	20121	$us uc cb$	1	1	-1	$us b + uc c, us c + uc b$	
	11121	$\frac{1}{\sqrt{2}}(us dc + ds uc) cb$	1	0	-1	$us b + dc c, us c + dc b$	-1.84
	02121	$ds dc cb$	1	-1	-1	$ds b + dc c, ds c + dc b$	
	11121	$ud sc cb$	0	0	-1	$ud b + sc c, ud c + sc b$	-1.84

Table 7. $[qqqqQ; Q = c (H_c)]$ The possible decay modes of H_c -dibaryons with configuration $qqqqQ$ ($Q = c$) against two baryon state B and B' . The numbers in the brackets are invariant mass of the two baryons at rest frame.

flavor	config. of H_c	I	I_z	S	config. of $B + B'$ (MeV)	final states
31110	$udusuc$	1	1	-1	$\Lambda_c^+ + \Sigma^+$ (3475.83)	$pK^-\pi^+ + p\pi^0$
					$p + \Xi_c^+$ (3406.17)	$p + \Lambda K^-\pi^+\pi^+$ $p + \Sigma^+ K^-\pi^+$ $p + \Xi^0\pi^+$ $p + \Xi^-\pi^+\pi^+$ $p + \Xi^0\pi^+\pi^0$ $p + \Xi^0\pi^+\pi^+\pi^-$
22110	$ud \frac{1}{\sqrt{2}}(usdc + dsuc)$	1	0	-1	$\Lambda + \Sigma_c^+$ (3568.58)	$p\pi^- + \Lambda_c^+\pi^0$
					$\Sigma^0 + \Lambda_c^+$ (3478.92)	$\Lambda\gamma + pK^-\pi^+$
13110	$ud dsdc$	1	-1	-1	$\Lambda_c^+ + \Sigma^-$ (3483.91)	$pK^-\pi^+ + n\pi^-$
					$n + \Xi_c^0$ (3410.56)	$n + pK^-K^-\pi^+$ $n + \Lambda K^-\pi^+\pi^+\pi^-$ $n + \Xi^-\pi^+$ $n + \Xi^-\pi^+\pi^+\pi^-$ $n + \Omega^- K^+$
21210	$udus sc$	1/2	1/2	-2	$\Lambda_c^+ + \Xi^0$ (3601.32)	$pK^-\pi^+ + \Lambda\pi^0$
					$\Lambda + \Xi_c^+$ (3583.58)	$p\pi^- + \Lambda K^-\pi^+\pi^+$ $p\pi^- + \Sigma^+ K^-\pi^+$ $p\pi^- + \Xi^0\pi^+$ $p\pi^- + \Xi^-\pi^+\pi^+$ $p\pi^- + \Xi^0\pi^+\pi^0$ $p\pi^- + \Xi^0\pi^+\pi^+\pi^-$
12210	$ud ds sc$	1/2	-1/2	-2	$\Lambda_c^+ + \Xi^-$ (3608.17)	$pK^-\pi^+ + \Lambda\pi^-$
					$\Lambda + \Xi_c^0$ (3586.68)	$p\pi^- + pK^-K^-\pi^+$ $p\pi^- + \Lambda K^-\pi^+\pi^+\pi^-$ $p\pi^- + \Xi^-\pi^+$ $p\pi^- + \Xi^-\pi^+\pi^+\pi^-$ $p\pi^- + \Omega^- K^+$
11310	$us ds sc$	0	0	-3	$\Xi^0 + \Xi_c^0$ (3785.86)	$\Lambda\pi^0 + pK^-K^-\pi^+$ $\Lambda\pi^0 + \Lambda K^-\pi^+\pi^+\pi^-$ $\Lambda\pi^0 + \Xi^-\pi^+$ $\Lambda\pi^0 + \Xi^-\pi^+\pi^+\pi^-$ $\Lambda\pi^0 + \Omega^- K^+$
					$\Xi^- + \Xi_c^+$ (3789.6)	$\Lambda\pi^- + \Lambda K^-\pi^+\pi^+$ $\Lambda\pi^- + \Sigma^+ K^-\pi^+$ $\Lambda\pi^- + \Xi^0\pi^+$ $\Lambda\pi^- + \Xi^-\pi^+\pi^+$ $\Lambda\pi^- + \Xi^0\pi^+\pi^0$ $\Lambda\pi^- + \Xi^0\pi^+\pi^+\pi^-$

Table 8. $[qqqqQ; Q = b (H_b)]$ The possible decay modes of H_b -dibaryons with configuration $qqqqQ$ ($Q = b$) against two baryon state B and B' . The numbers in the brackets are invariant mass of the two baryons at rest frame.

flavor	config. of H_b	I	I_z	S	config. of $B + B'$ (MeV)	final states
31101	$ud us ub$	1	1	-1	$p + \Xi_b^0$ (6730.67)	$p + ?$
					$\Lambda_b^0 + \Sigma^+$ (6809.57)	$\Lambda_c^+ \pi^- + p \pi^0$
22101	$ud \frac{1}{\sqrt{2}}(us db + ds ub)$	1	0	-1	$\Lambda + \Sigma_b^0$ (6926.18)	$p \pi^- + \Lambda_b^0 \pi^0$
					$\Sigma^0 + \Lambda_b^0$ (6812.84)	$\Lambda \gamma + \Lambda_c^+ \pi^-$
13101	$ud ds db$	1	-1	-1	$\Lambda_b^0 + \Sigma^-$ (6817.0)	$\Lambda_c^+ \pi^- + n \pi^-$
					$n + \Xi_b^-$ (6731.97)	$n + ?$
21202	$ud us sb$	1/2	1/2	-2	$\Lambda_b^0 + \Xi^0$ (6935.06)	$\Lambda_c^+ \pi^- + \Lambda \pi^0$
					$\Lambda + \Xi_b^0$ (6908.08)	$p \pi^- + ?$
12201	$ud ds sb$	1/2	-1/2	-2	$\Lambda_b^0 + \Xi^-$ (6941.9)	$\Lambda_c^+ \pi^- + \Lambda \pi^-$
					$\Lambda + \Xi_b^-$ (6908.08)	$p \pi^- + ?$
11301	$us ds sb$	0	0	-3	$\Xi_b^0 + \Xi^-$ (7114.1)	$? + \Lambda \pi^-$
					$\Xi^0 + \Xi_b^-$ (7107.26)	$\Lambda \pi^0 + ?$

Table 9. $[qqqQqQ'; Q = c, Q' = b (H_{bc})]$ The possible decay modes of H_{bc} -dibaryons with configuration $qqqQqQ'$ ($Q = c, Q' = b$) against two baryon state B and B' . The numbers in the brackets are invariant mass of the two baryons at rest frame.

flavor	config. of H_{bc}	I	I_z	S	config. of $B + B'$ (MeV)	final states
31011	$ud uc ub$	1	1	0	$\Lambda_b^0 + \Sigma_c^{++}$ (8074.2)	$\Lambda_c^+ \pi^- + \Lambda_c^+ \pi^+$
					$p + \Xi_{cb}^+$ (?)	$p + ?$
22011	$ud \frac{1}{\sqrt{2}}(uc db + dc ub)$	1	0	0	$\Xi_{cb}^0 + p$ (?)	$? + p$
					$n + \Xi_{cb}^+$ (?)	$n + ?$
13011	$ud dc db$	1	-1	0	$\Lambda_b^0 + \Sigma_c^0$ (8073.96)	$\Lambda_c^+ \pi^- + \Lambda_c^+ \pi^-$
					$n + \Xi_{cb}^0$ (?)	$n + ?$
30111	$us uc ub$	3/2	3/2	-1	$\Sigma^+ + \Xi_{cb}^+$ (?)	$p \pi^0 + ?$
					$\Xi_b^0 + \Sigma_c^{++}$ (8246.4)	$? + \Lambda_c^+ \pi^+$
21111	$\frac{1}{\sqrt{3}}(us uc db + us dc ub + ds uc ub)$	3/2	1/2	-1	$\Sigma_b^0 + \Xi_c^+$ (8278.4)	$\Lambda_b^0 \pi^0 + \Lambda K^- \pi^+ \pi^+$ $\Lambda_b^0 \pi^0 + \Sigma^+ K^- \pi^+$ $\Lambda_b^0 \pi^0 + \Xi^0 \pi^+$ $\Lambda_b^0 \pi^0 + \Xi^- \pi^+ \pi^+$ $\Lambda_b^0 \pi^0 + \Xi^0 \pi^+ \pi^0$ $\Lambda_b^0 \pi^0 + \Xi^0 \pi^+ \pi^+ \pi^-$
					$\Sigma_c^+ + \Xi_b^0$ (8245.3)	$\Lambda_c^+ \pi^0 + ?$
12111	$\frac{1}{\sqrt{3}}(us dc db + ds dc ub + ds uc db)$	3/2	-1/2	-1	$\Sigma_b^0 + \Xi_c^0$ (8282.5)	$\Lambda_b^0 \pi^0 + p K^- K^- \pi^+$ $\Lambda_b^0 \pi^0 + \Lambda K^- \pi^+ \pi^+ \pi^-$ $\Lambda_b^0 \pi^0 + \Xi^- \pi^+$ $\Lambda_b^0 \pi^0 + \Xi^- \pi^+ \pi^+ \pi^-$ $\Lambda_b^0 \pi^0 + \Omega^- K^+$
					$\Sigma_c^+ + \Xi_b^-$ (8309.9)	$\Lambda_c^+ \pi^0 + ?$
03111	$ds dc db$	3/2	-3/2	-1	$\Xi_b^- + \Sigma_c^0$ (8246.16)	$? + \Lambda_c^+ \pi^-$
					$\Sigma^- + \Xi_{cb}^0$ (?)	$n \pi^- + ?$
10311	$us sc sb$	1/2	1/2	-3	$\Xi_b^0 + \Omega_c^0$ (8489.9)	$? + \Sigma^+ K^- K^- \pi^+$ $? + \Xi^0 K^- \pi^+$ $? + \Xi^- K^- \pi^+ \pi^+$ $? + \Omega^- \pi^+$ $? + \Omega^- \pi^+ \pi^0$ $? + \Omega^- \pi^+ \pi^+ \pi^-$
					$\Xi^0 + \Omega_{cb}^0$ (?)	$\Lambda \pi^0 + ?$
01311	$ds sc sb$	1/2	-1/2	-3	$\Xi_b^- + \Omega_c^0$ (8489.9)	$? + \Sigma^+ K^- K^- \pi^+$ $? + \Xi^0 K^- \pi^+$ $? + \Xi^- K^- \pi^+ \pi^+$ $? + \Omega^- \pi^+$ $? + \Omega^- \pi^+ \pi^0$ $? + \Omega^- \pi^+ \pi^+ \pi^-$
					$\Xi^- + \Omega_{cb}^0$ (?)	$\Lambda \pi^- + ?$

Table 10. $[qq qq QQ'; Q = c, Q' = b (H'_{bc})]$ The possible decay modes of H'_{bc} -dibaryons with configuration $qq qq QQ'$ ($Q = c, Q' = b$) against two baryon state B and B' . The numbers in the brackets are invariant mass of the two baryons at rest frame.

flavor	config. of H'_{bc}	I	I_z	S	config. of $B + B'$ (MeV)	final states
21111	$ud us cb$	1/2	1/2	-1	$\Lambda_b^0 + \Xi_c^+$ (8088.1)	$\Lambda_c^+ \pi^- + \Lambda K^- \pi^+ \pi^+$ $\Lambda_c^+ \pi^- + \Sigma^+ K^- \pi^+$ $\Lambda_c^+ \pi^- + \Xi^0 \pi^+$ $\Lambda_c^+ \pi^- + \Xi^- \pi^+ \pi^+$ $\Lambda_c^+ \pi^- + \Xi^0 \pi^+ \pi^0$ $\Lambda_c^+ \pi^- + \Xi^0 \pi^+ \pi^-$
					$\Lambda_c^+ + \Xi_b^0$ (8078.86)	$p K^- \pi^+ + ?$
12111	$ud ds cb$	1/2	-1/2	-1	$\Lambda_b^0 + \Xi_c^0$ (8091.2)	$\Lambda_c^+ \pi^- + p K^- K^- \pi^+$ $\Lambda_c^+ \pi^- + \Lambda K^- \pi^+ \pi^+ \pi^-$ $\Lambda_c^+ \pi^- + \Xi^- \pi^+$ $\Lambda_c^+ \pi^- + \Xi^- \pi^+ \pi^+ \pi^-$ $\Lambda_c^+ \pi^- + \Omega^- K^+$
					$\Lambda_c^+ + \Xi_b^-$ (8078.86)	$p K^- \pi^+ + ?$
11211	$us ds cb$	0	0	-2	$\Xi_b^+ + \Xi_c^0$ (8263.4)	$? + p K^- K^- \pi^+$ $? + \Lambda K^- \pi^+ \pi^+ \pi^-$ $? + \Xi^- \pi^+$ $? + \Xi^- \pi^+ \pi^+ \pi^-$ $? + \Omega^- K^+$
					$\Xi_c^+ + \Xi_b^0$ (8260.3)	$\Lambda K^- \pi^+ \pi^+ + ?$ $\Sigma^+ K^- \pi^+ + ?$ $\Xi^0 \pi^+ + ?$ $\Xi^- \pi^+ \pi^+ + ?$ $\Xi^0 \pi^+ \pi^0 + ?$ $\Xi^0 \pi^+ \pi^+ \pi^- + ?$

Table 11. $[qq qQ QQ'; Q = c, Q' = b (H_{bcc})]$ The possible decay modes of H_{bcc} -dibaryons with configuration $qq qQ QQ'$ ($Q = c, Q' = b$) against two baryon state B and B' . The numbers in the brackets are invariant mass of the two baryons at rest frame.

flavor	config. of H_{bcc}	I	I_z	S	config. of $B + B'$ (MeV)	final states
21021	$ud uc cb$	1/2	1/2	0	$\Lambda_b^0 + \Omega_{cc}^+$ (?)	$\Lambda_c^+ \pi^- + ?$
					$\Lambda_c^+ + \Xi_{cb}^+$ (?)	$p K^- \pi^+ + ?$
12021	$ud dc cb$	1/2	-1/2	0	$\Lambda_b^0 + \Omega_{cc}^0$ (?)	$\Lambda_c^+ \pi^- + ?$
					$\Lambda_c^+ + \Xi_{cb}^0$ (?)	$p K^- \pi^+ + ?$
20121	$us uc cb$	1	1	-1	$\Xi_b^+ + \Omega_{cc}^+$ (?)	$? + ?$
					$\Lambda_c^+ + \Xi_{cb}^+$ (?)	$\Lambda K^- \pi^+ \pi^+ + ?$ $\Sigma^+ K^- \pi^+ + ?$ $\Xi^0 \pi^+ + ?$ $\Xi^- \pi^+ \pi^+ + ?$ $\Xi^0 \pi^+ \pi^0 + ?$ $\Xi^0 \pi^+ \pi^+ \pi^- + ?$
11121	$\frac{1}{\sqrt{2}}(us dc + ds uc) cb$	1	0	-1	$\Lambda_b^0 + \Omega_{cc}^{++}$ (?)	$\Lambda_c^+ \pi^- + ?$
					$\Lambda_c^+ + \Omega_{cb}^0$ (?)	$p K^- \pi^+ + ?$
02121	$ds dc cb$	1	-1	-1	$\Xi_b^0 + \Xi_{cc}^+$ (?)	$? + ?$
					$\Xi_c^0 + \Xi_{cb}^0$ (?)	$p K^- K^- \pi^+ + ?$ $\Lambda K^- \pi^+ \pi^+ \pi^- + ?$ $\Xi^- \pi^+ + ?$ $\Xi^- \pi^+ \pi^+ \pi^- + ?$ $\Omega^- K^+ + ?$
11121	$ud sc cb$	0	0	-1	$\Lambda_b^0 + \Omega_{cc}^+$ (?)	$\Lambda_c^+ \pi^- + ?$
					$\Lambda_c^+ + \Xi_{cb}^+$ (?)	$p K^- K^- \pi^+ + ?$